

Full Length Research Paper

Interaction effects of in-organic N-fertilizer and seed rates on yield and quality traits of malt barley varieties in the Highland of Tigray, North Ethiopia

Yemane Nega^{1*} Addis Abraha² and Girmay G. Samuel³

¹Tigray Agricultural Research Institute, Alamata Research Center, P.O. Box 56, Alamata, Ethiopia.

²Mekelle University, Department of Dryland Crop and Horticultural Sciences, P.O. BOX 231, Ethiopia.

³Mekelle University, Department of Natural Resources Management, P.O. BOX 231, Ethiopia.

Accepted 27 July, 2015

A new brewery factory has been constructed in Tigray region. In response, farmers' concerns are to produce and earn premium price from malt barley. Besides, quality prerequisites for malting barley are objectively strict and mostly controlled by the growers' crop management practice. However, there is limited/no information on the fertilizer and seed rate requirement of the crop that can achieve high quality malt without compromising yield and profit. Therefore, this research was aimed to determine the appropriate levels of inorganic N-fertilizer and seed rate on the yield and quality of malt barley varieties in three districts of Southern Tigray, North of Ethiopia. Treatments include two malt barley varieties (V1=Bekoji-1 and V2=Bhatty), seven N-rates (N0=0, N1=23, N2=34, N3=46, N4=57.5, N5=69 and N6=80.5 Kg N ha⁻¹) in the form of urea and two levels of seed rates (SR1=100, SR2=150 kg ha⁻¹). The design was split-split plot with three replications. Varieties were put as a main plot, seed rate as a sub plot and the N-rates as a sub sub-plot treatment. The analysis of the separate and combined over locations revealed significant differences on grain yield (GY) and malt quality traits due to the main effect of treatments and some interactions. The effect of variety and seed rate were non-significant in response to GY. Highest value of grain yield (GY), hectoliter weight, plumpness, thousand kernel weight and acceptable grain protein content (GP) for malt barley was recorded at N5. The treatment without N-fertilizer application showed the least performance in above ground biomass and grain yield and malt quality traits. From the interaction of factors acceptable, GP and higher grain yields were obtained from combinations V2×SR1×N5 (GP=11.22%) and V1×SR1×N5 (GP=12.50%) at Ofla, combinations V2×SR1×N6 (GP=12.28%) and V2×SR1×N5 (GP=12.09%) at Enda-Mekoni, and combinations V1×SR1×N5 (GP=11.94%) and V2×SR1×N5 (GP=11.44%) at Emba-Alaje. Combination V2×SR1×N5 was selected as the best combination in all locations. Bekoji-1(V1) at higher N-rates (N5 and N6) and SR1 at Enda-Mekoni were beyond the maximum acceptable limit in grain protein content. Regarding the profitability of the package, the highest net benefit and MRR were obtained from the application of N5 fertilizer rate at SR1 for both varieties in all study areas. Therefore, under similar environmental conditions, the combination of V2×SR1×N5 is the best agronomic package for both grain yield and acceptable malt quality at all locations. In addition, combinations of V1×SR1×N5 can also be recommended for Ofla and Emba-Alaje districts.

Key words: Grain protein, grain yield, N-fertilizer, profit, seed rate, variety.

INTRODUCTION

Barley (*Hordeum vulgare* L.) is the fourth important cereal crop in the world and fifth important cereal in Ethiopia (Bayeh and Birhane, 2006). It is largely grown as a food crop in the central and northern parts of Ethiopia, with Oromia, Amhara, Tigray, and Southern Nations, Nationalities, and People's Region (SNPPR) as the main

areas of production (ATA, 2012). The use of malt barley as a raw material in brewery factory has increased its

*Corresponding author. E-mail: yema14@gmail.com.

value and the demand of farmers to produce (ATA, 2012). Barley malt is the perfect combination of starch, enzymes, flavors, and aromas for brewing. There are many types of barley malt – from light to dark but all are variations on two principal themes: germination and kilning. Different end-uses require different malt quality specifications. Some of the principal characteristics used to define malting quality are protein (low, moderate, or high), malt extract (high), enzyme activity (moderate to high), and beta glucan (low).

Despite the immense potential for producing malt barley in Ethiopia, only about 2% of total barley produced goes into malt factory for the six local breweries in the country (Tefera, 2012). Only one-third can be supplied from locally produced barley. The remaining two-thirds are imported primarily from Belgium and France (ATA, 2012). Given the quantity needed, by brewery factories in Ethiopia [BGI-Castel, Diageo (Meta, Zemen)], Dashen, Bedele, Heineken [(Harar and Walya) and Raya beer] demand for malt barley is expected to reach 110,000 to 130,000 MT in the year 2016. The reason for the low production of malt barley in domestic market is mainly due to the emphasis given to limited areas. The Arsi and Bale highland has been the only areas chosen for the production of malt barley for many years. This limits the expansions in area coverage to other potential areas. Therefore, it is crucial and timely to expand the malt barley production in different barley growing regions of the country to satisfy the ever-growing demand for raw materials by the beverage industry and reduce the volume of imported malt barley.

Regardless of the available conducive environment in the country, there have been limited researches done to evaluate the yield and malt quality traits of the barley varieties in response to different rates of nitrogen fertilizer and agronomic practices. The most important limiting factors which highly affect the yield and the malt quality of barley are variety, fertilization, seed rate, sowing time, and soil moisture (Grant, 2000). Nitrogen (N) fertilizer is the main input necessary for production of high quality malt barley. Therefore, the objective of this study was to investigate response of two varieties of malt barley to different rates of nitrogen fertilizer (N) and seed rates across three locations in the highland of Southern Tigray, Northern Ethiopia.

MATERIALS AND METHODS

Site description

Field experiment was conducted to investigate response of two malt varieties to different rates of nitrogen fertilizer (N) and seed rates across three locations. The experiment was conducted in 2013 growing season under rain fed conditions. The experimental fields were located in Ofla district at Awlie-Gara research station, Emba-Alaje districts at Ayba farmers' training center and

Enda-Mehoni district at Mekan on farmers' land. The geographic location and agro-ecological condition of the study sites is mentioned in Table 1 and Figure 1.

Soil physical and chemical properties

Soil texture, pH, total N, available P, cation exchange capacity, organic carbon and electrical conductivity were determined for the composited soil samples collected from the experimental field at 0-30 cm depth before sowing as indicated in Table 2.

Treatments and experimental design

The experiment includes two malt barley varieties (V1=Bekoji-1 and V2=Bhatty), seven nitrogen fertilizer rates (N0=0, N1=23, N2=34, N3=46, N4=57.5, N5=69 and N6=80.5 Kg N ha⁻¹) and two levels of seed rates (SR1=100, SR2=150 kg ha⁻¹). The design was a split-split plot with three replications. Varieties were put as a main plot, seed rate as a sub plot and the N-rates as a sub sub-plot treatment. Plot size was 2*2.5m² (10 rows), and harvestable plot size was 1.6*2.5m² (8 rows). Spacing between rows, sub-sub-plots, sub-plots, main plots and replications was 0.2, 1, 1, 1.0 and 1.5 m, respectively. Nitrogen in the form of urea (46%N) was applied in three splits, that is, at planting, at first weeding and anthesis stage and TSP-46% P₂O₅ as phosphorus source. Phosphorus was applied uniformly for all treatments at the rate of 46 P₂O₅ kg ha⁻¹.

Data collection

Agronomic data were collected on grain yield and biological yield. In addition malt quality trait data were collected on thousand kernel weights, hectoliter weight, sieve test/kernel plumpness and grain protein content. Hectoliter weight was determined using a standard laboratory hectoliter weight apparatus (EASY-WAY hectoliter weight test machine) and electronic balance according to the AACC (2000) Method 55-10, and the obtained values were adjusted to a moisture content of 12.5% basis. Sieve test/kernel plumpness showed that barley grain sample (100 g) from each plot was sieved on 2.8, 2.5, and 2.2 mm slotted sieve. The mass left on each sieve size was measured and the percentage was computed as described in the European Brewery Convention (EBC, 1987). Total grain protein content was determined by multiplying Grain N-content x 6.25. Grain N-content was determined by micro-Kjeldahl method as percentage of dry matter as stated in the ESA method ES 669-2001(ESA, 2001).

Statistical analysis

All data were presented as mean values of three replicates. Data were analyzed statistically following the

Table 1. Area description of the experiment sites.

Study site	Lat. (N)	Long. (E)	Alt. (masl)	Mean RF (mm)		Min T (°C)	Max. T (°C)
				Jun-Sept	Oct-May		
Ofla	12° 31'	39° 33'	2546	654.42	345.96	5.4	20.2
Endamekoni	12° 44'	39° 32'	2423	462.05	250.31	10.3	22.6
Emba-Alaje	12° 54'	39° 31'	2724	518.55	67.28	10.3	22.6




Figure 1. Monthly rain fall distribution of the research sites during the study (2013).
Source: National Meteorological Agency.

Table 2. Major soil properties of the experimental sites before planting.

Soil properties	Location			Method
	Ofla	Enda-Mekoni	Emba-Alaje	
pH	6.167 ^{sa}	6.538 ^{sa}	6.155 ^{sa}	pH meter (1:25)
OM (%)	2.343 ^L	2.652 ^L	3.473 ^{Opt}	Walkley and Black
OC (%)	1.359 ^L	1.539 ^L	2.014 ^{Opt}	Walkley and Black
Total N (%)	0.1372 ^{VL}	0.133 ^{VL}	0.1946 ^{VL}	Kjeldahl
C:N	9.91 ^m	11.57 ^m	10.35 ^m	%TN:%OC
P (ppm)	13.6 ^{VL}	10.32 ^{VL}	35.96 ^{Opt}	Olisen II
CEC	19.2 ^m	36.4 ^h	31.4 ^h	Amm. Acet extraction
ECe (mS/cm) (1:2.5)	0.17 ^{ns}	0.16 ^{ns}	0.16 ^{ns}	ESSP
Soil texture	Clay loam	Loam	Clay loam	Sieve test
Soil type	Vertisol	Vertisol	Vertisol	-

h = high, m = medium, vh = very high, L = low, vl = very low, sa = slightly acidic, N = Neutral, ns = non-salin.

Source: Ethio SIS team analysis (2013).

method described by Gomez and Gomez (1984). Analysis of variance (ANOVA) was carried out using Genstat (13th edition) computer software package to determine the level of significance at 95%. Fisher's protected LSD test at 5% probability was used for the mean comparison when the analysis of variance indicated the presence of significant differences.

RESULTS AND DISCUSSION

Soil test results

According to the fertility classification of Landon (1991),

the soil analysis of the locations shows that the soils of the experimental sites were vertisol, slightly acidic, low to medium content of organic carbon (OC), very low content of available P and total N content, medium to high CEC and nono salin soil for crop growth (Table 2).

Biomass and grain yield of malt barley

The analyses of variance showed that varieties do not exhibit significant difference on the above ground biomass except in Emba-Alaje. Comparisons of treatment means at Emba-Alaje indicated relatively that higher above ground biomass yield was recorded from variety

Table 3. The main effects of N-rate, seed rate and variety on biomass and grain yield.

Treatment	Biomass (kg ha ⁻¹)				Grain yield (kg ha ⁻¹)			
	Ofla	Enda- Mekoni	Emba-Alaje	Mean	Ofla	Enda- Mekoni	Emba-Alaje	Mean
N-Rates (kg ha⁻¹)								
0	8850 ^f	8990 ^f	10900 ^g	9580 ^g	1874 ^d	1966 ^e	3598 ^e	2479 ^e
23	10590 ^e	10040 ^e	12260 ^f	10960 ^f	2289 ^c	2354 ^d	4451 ^d	3031 ^d
34.5	11050 ^e	11100 ^d	13160 ^e	11770 ^e	2536 ^c	2671 ^c	4708 ^d	3305 ^c
46	12060 ^d	12030 ^c	14000 ^d	12690 ^d	2930 ^b	3052 ^b	5052 ^c	3678 ^b
57.5	13210 ^c	13130 ^b	14770 ^c	13700 ^c	3001 ^{ab}	3169 ^b	5207 ^{bc}	3793 ^b
69	14260 ^b	14890 ^a	15840 ^b	15000 ^b	3322 ^a	3490 ^a	5615 ^a	4142 ^a
80.5	15430 ^a	15280 ^a	17040 ^a	15920 ^a	3231 ^{ab}	3534 ^a	5352 ^{ab}	4039 ^a
SE (M)±	341.4	238.3	169.6	232.8	170	124.7	134.6	97.1
LSD 0.05	686.5	479.1	341.0	459.6	341.8	250.8	270.7	191.8
Seed rates (kg ha⁻¹)								
100	11435	12292	13994	12574	2596	2907	4833	3445.89
150	12980	12124	14001	13035	2870	2875	4877	3544.78
SE (M)±	508.4	60.5	164.2	124.4	111.9	113.4	64.6	51.9
LSD 0.05	1411.5	167.9	ns	245.7	ns	ns	ns	ns
Variety								
Bekoji-1	12132	12357	13798	12762	2611	2866	4878	3451.72
Bhatty	12284	12059	14197	12846	2870	2916	4831	3538.95
SE (M)±	651.2	192.1	61.5	124.4	95.7	29.8	134.5	51.9
LSD 0.05	ns	ns	264.7	ns	ns	ns	ns	ns
CV	690	480	3	770	15.2	10.6	6.8	11.7
GM	12208	12208	13997	12804	2740	2891	4855	3504.86

ns = non-significant, SE (M) ± = standard error of mean, LSD = least significant difference, CV = coefficient of variation, GM = grand mean.

Columns not connected by similar letters have significant difference.

Bhatty (14197 vs. 13798 kg ha⁻¹) (Table 3). The effect of seed rate on the above ground biomass was found to be significant at Ofla ($P < 0.05$) and Enda-Mekoni ($P < 0.01$) but not at Emba-Alaje. Highest biomass was recorded from the highest seed rate at Ofla (12980 vs. 11435 kg ha⁻¹). In contrast, highest biomass was recorded from the lower seed rate at Enda-Mekoni (12292 vs. 12124 kg ha⁻¹) (Table 3). This shows that response of biomass yield to seed rate varied across locations and generally it can be lower or higher depending on the soil moisture content and fertility status of the experimental fields. Grain yield response was non-significant due to variation of seed rate and variety. The effect of seed rate has been documented previously in barley (Jedel and Helm, 1995), in wheat (Stougaard and Xue, 2004) and malt barley (O'Donovan et al., 2011, 2013) that under similar environments, variety and seed rate have non-significant effect on grain yield.

Analysis of variance showed that N-fertilizer rate was significantly ($P \leq 0.001$) in response to biological and grain yield of malt barley (Table 3) at all locations. Comparison of treatment means combined from three locations

indicated that biological yield generally increased with the increase in the rate of N-fertilizer. Hence, maximum biological yield was obtained from 80.5 kg N ha⁻¹, whereas the lowest biomass yield was obtained from plants grown without N-fertilizer (Table 3). Plants supplied with the highest rate of N had biological yields exceeding 66.5% from plants in the control treatment (Table 3). The result was in line with that of Alam et al. (2005) who reported that accumulation of dry matter of barley increased with higher doses of N-fertilizer rate. Malt barley grain yield generally increased with an increase in rate of nitrogen up to 69 kg ha⁻¹. Hence, the highest grain yield at Ofla (3322 kg ha⁻¹) and Emba-Alaje (5615 kg ha⁻¹) was obtained from 69 kg ha⁻¹. In the case of Enda-Mekoni, the highest grain yield (3534 kg ha⁻¹) was recorded at the maximum rate of N-fertilizer (N6). Lowest grain yield in all experimental fields was obtained from the control (N0) (Table 3). The grain yield obtained from N5 exceeded by an additional 67% increment of the one obtained from the control treatment (Table 3) indicates that deficiency of N is evident in the reduction of light interception by decreasing leaf area index, which

Table 4. Effect of variety and seed rate interaction on biomass and grain yield.

Treatment		Biomass (Q ha ⁻¹)				Grain yield (kg ha ⁻¹)			
Seed rate (kg ha ⁻¹)	Variety	Ofla	Enda- Mekoni	Emba-Alaje	Mean	Ofla	Enda- Mekoni	Emba-Alaje	Mean
100	Bekoji-1	11040 ^c	12180 ^a	13610 ^c	12280 ^c	2630 ^b	2851	4779 ^{abc}	3410.27
	Bhatty	11830 ^b	12400 ^a	14380 ^a	12870 ^b	2624 ^b	2963	4886 ^{ab}	3481.52
150	Bekoji-1	13220 ^a	12530 ^a	13990 ^b	13250 ^a	2650 ^b	2881	4977 ^a	3493.18
	Bhatty	12740 ^a	11720 ^b	14020 ^b	12820 ^b	3173 ^a	2869	4776 ^{bc}	3596.38
SE (M)±		826.2	2.014	137.4	176.0	131.2	117.3	105.7	73.421
LSD		2219.3	7.407	275.5	347.4	263	ns	211.8	ns
CV %		9.5	4.8	3.2	7.7	15.4	10.6	7.1	11.8

ns = non-significant, SE (M) ± = standard error of mean, LSD = least significant difference, CV = coefficient of variation, GM = grand mean. Columns not connected by similar letters have significant difference.

Table 5. Interaction of variety, seed rate and N-rate on grain yield.

N-rate (kg ha ⁻¹)	Seed rate (kg ha ⁻¹)	Grain yield (kg ha ⁻¹)							
		Ofla		Enda-Mekoni		Emba-Alaje		Mean	
		Var-1	Var-2	Var-1	Var-2	Var-1	Var-2	Var-1	Var-2
0	100	1628	2049	2010	2183	3785	3396	2474	2543
	150	1737	2080	1910	1762	3488	3721	2378	2521
23	100	1987	2245	2285	2362	4462	4447	2911	3018
	150	2103	2820	2455	2315	4336	4560	2965	3232
34.5	100	2412	2348	2492	2723	4634	4686	3179	3253
	150	2435	2950	2598	2872	4740	4772	3258	3531
46	100	2808	2612	2992	2882	4833	5113	3544	3536
	150	2913	3386	3162	3175	5053	5209	3709	3923
57.5	100	3008	2905	3128	3190	5062	5199	3733	3765
	150	2644	3447	3182	3177	5660	4908	3829	3844
69	100	3262	3043	3442	3623	5503	5813	4069	4160
	150	3322	3663	3323	3570	5876	5266	4174	4167
80.5	100	3101	2965	3607	3777	5175	5551	3961	4098
	150	3195	3662	3538	3213	5687	4996	4140	3957
GM		2611	2870	2866	2916	4878	4831	3452	3539
V×SR×NR									
LSD 0.05		Ns	Ns	Ns	560.5	ns			
SE (M)±		347.1	262.9	279.6	194				
CV		15.5	11.1	7.1	11.8				

Var-1 = Bekoji-1 and Var-2 = Bhatty.

results in lower grain yield. This result was in support of that of many authors namely: Moreno et al. (2003), Alam et al. (2005) and Shafi et al. (2011) in barley. Similarly Shafi et al. (2011) also reported that nitrogen applied at the rate of 60 kg ha⁻¹ resulted in maximum grain spike⁻¹, thousand grain weight, biological yield, grain yield, and grain and plant N-content. Conversely, grain yield was reduced by 43.6, 44.4 and 35% from the highest recorded in the absence of N-fertilizer in the control at Ofla, Enda-Mekoni and Emba-Alaje, respectively (Table 4). This was in support of Stewart (2002) who reviewed

that approximately 41% (in corn), 27% (in rice), 19% (in barley and sorghum), 16% (in wheat) yield reduction was contributed by eliminating N-fertilizer on the above mentioned crop yields. The interaction of variety to seed rate was significant in response to BM and GY. At higher seed rate, variety Bekoji-1 achieves more grain yield and vice versa. Moreover, the third level interaction of variety, seed rate and N-fertilizer rate was significant in response of GY at Emba-Alaje. The highest average GY was recorded from the interaction of V1×SR2×N5 and V2×SR1×N5 (Table 5). The result of this study confirmed

Table 6. Analysis of variance for TKW, grain protein and HLW of malt barley at each location.

Source of variation	Mean squares for source of variation								
	TKW (gm)			% GPC			HLW		
	Ofla	E/Mekoni	E/Alaje	Ofla	E/Mekoni	E/Alaje	Ofla	Mekan	E/Alaje
Rep (1)	36.0833	0.9048	0.7262	0.0444	4.3055	2.6839	13.26	1.51143	0.05786
NR (6)	0.0476 ^{ns}	0.7619 ^{ns}	0.7619 ^{ns}	4.2258 ^{***}	5.2999 ^{**} *	3.8996 ^{**} *	2.7304 ^{**} *	3.8795 ^{**} *	4.9307 ^{***}
SR (1)	2.5833	0.7619	0.5119	0.7622+	1.9515 ^{**}	0.0279 ^{ns}	1.1286 ^{**}	0.02571 ⁿ _s	0.52071 ^{**} *
Var (1)	40.0476 [*] *	48.7619 ^{**} *	1.1905+	21.7077 ^{**} *	5.2361 ^{**} *	4.316 ^{***}	1.8036 ^{**} *	0.18286 [*] *	0.20643 ^{**}
NR×SR (6)	0.0476 ^{ns}	5.7619 ^{**}	0.0476 ^{ns}	0.077 ^{ns}	0.0149 ^{ns}	0.1065 ^{ns}	0.1294 ^{ns}	0.03571 ⁿ _s	0.05738 [*]
NR×Var (6)	0.9762	0.5476	0.1905	0.1642 ^{ns}	0.064 ^{ns}	0.0216 ^{ns}	0.0694 ^{ns}	0.01952 ⁿ _s	0.0231 ^{ns}
SR×Var (1)	24.746 ^{**} *	27.0675 ^{**} *	40.6032 ^{**} *	0.3816 ^{ns}	0.0223 ^{ns}	0.0428 ^{ns}	0.01 ^{ns}	0.10286 [*]	0.315 ^{***}
NR×SR×Var (6)	0.9921 ^{ns}	0.5119 ^{ns}	0.0952 ^{ns}	0.0586 ^{ns}	0.0493 ^{ns}	0.0335 ^{ns}	0.1059 ^{ns}	0.01286 ⁿ _s	0.03833+
Residual (27)	2.6032 ^{**}	0.2897 ^{ns}	1.3016 ^{ns}	0.1868	0.1844	0.2692	0.1137	0.0181	0.01786

TKW = 1000 kernel weight, GPC = grain protein content, and HLW = hectoliter weight (kg ha⁻¹).

that the rate of N-fertilizer is the most influential factor in biomass and grain yield of malt barley. Similar results were reported in the studies of Ahmad (1999), Miao et al. (2006) and Oikeh et al. (2007) that nitrogen plays a very important role in crop productivity.

Malt quality traits

The main effect of N-rate, seed rate and variety were significant on most of the malt quality traits (Tables 6 and 7). In this study, the varieties had an acceptable range of grain protein concentration in reference to Canadian Grain Commission (CGC) and Ethiopian Standard Agency (ESA) standards except in some of the third level treatment interactions at Ofla and Enda-Mekoni. Variety Bekoji-1 was relatively higher in grain protein accumulation in all experimental fields (Table 6). The variation in grain protein of malt barley varieties grown side by side was in support of that of many authors (O'Donovan et al., 2011, 2013). Relatively higher grain protein (GP), hectoliter weight (HLW), kernel plumpness (KP) and thousand kernel weight (TKW) were recorded from 100 kg ha⁻¹ (SR1) (Table 7). Grain protein content, TKW, KP and HLW had a positive response to N-fertilizer rate and was higher at N5 fertilizer application, while low values were obtained from the control (N0) (Figure 2). Low value of TKW, KP and HLW indicates poor grain filling. Slight increases in specific weight in response to N-application have been reported by most authors (Pushman and Bingham, 1976; Gooding and Davies, 1997) under more favorable growing conditions (Figure

1). In addition, McKenzie and Jackson (2005) reported that an increase in N-fertilizer supply resulted in an increase in grain yield and grain protein. However, GP in all main effect of treatments was within the acceptable standard range for malt barley. This may be due to the genetic nature of the varieties for malt purpose. Emphasis should be given for the third level interaction of treatments as far as the quality standard is the objective. Interaction of V1×SR1×N6, (GP=12.72%) at Ofla, V1×SR1×N5 (GP=12.81), V1×SR1×N6 (GP=13) (GP=13.0), V1×SR2×N5 (GP=12.69) and V1×SR2×N6 (GP=12.75) at Enda-Mekoni were treatment combinations with grain protein content above maximum acceptable limit for malt barley (Table 8). According to the standards of CGC and ESA grain protein content, >12.5% is unacceptable for malt purpose. In general, Emba-Alaje was a suitable environment in response to grain protein for malt purpose while yield was enhanced. Moreover, variety Behatty has relatively low grain protein accumulation nature even at higher N-rate and seed rate.

Partial budget analysis

According to the partial budget analysis (PBA) technique of CIMMYT (1988) of N-fertilizer on malt barley, the highest net benefit was obtained from the application of 69 N5 kg N ha⁻¹ at all locations. However, the highest marginal rate of return (MRR) was recorded from 46 kg N ha⁻¹ (N3), and N5 at Ofla and Enda-Mekoni, while N5 at Emba-Alaje (Table 9). Finally, N5 for both varieties can be selected for its high net benefit and because it can

Table 7. The effect of variety and seed rate on grain protein, hectoliter weight and 1000 kernel weight of malt barley.

Treatment	Grain protein (%)				Hectoliter weight (kg HL ⁻¹)				1000 kernel weight (gm.)			
	Ofla	Enda-Mekoni	Emba-Alaje	Mean	Ofla	Enda-Mekoni	Emba-Alaje	Mean	Ofla	Enda-Mekoni	Emba-Alaje	Mean
Seed rate (kg ha⁻¹)												
100	11.358	11.577	11.18	11.37	61.9	62.7	63.2	62.585	44.24	45.02	45.64	44.97
150	11.125	11.203	11.135	11.16	61.6	62.7	63.0	62.412	42.86	43.50	45.41	43.92
SE (M)±	0.1155	0.1148	0.1387	0.0835	0.0901	0.036	0.0357	0.054	0.2156	0.1615	0.0952	0.1203
LSD	ns	0.24	ns	0.17	0.19	ns	0.07	0.11	0.6	0.45	ns	0.24
Variety												
Bekoji-1	11.864	11.696	11.435	11.05	61.6	62.8	63.1	62.478	43.57	44.36	45.62	44.52
Bhatty	10.619	11.084	10.88	11.48	61.9	62.7	63.0	62.52	43.52	44.17	45.43	44.37
SE (M)±	0.1155	0.1148	0.1387	0.0835	0.0901	0.036	0.0357	0.054	0.3507	0.1905	0.1561	0.1203
LSD 0.05	0.24	0.24	0.29	0.17	0.19	0.07	0.07	0.11	1.51	ns	ns	ns
CV	3.8	3.8	4.7	4.8	0.5	0.2	0.2	0.6	2.0	1.7	1.9	2.1
GM	11.24	11.39	11.16	11.26	61.7	62.7	63.1	62.5	43.55	44.26	45.52	44.44




Figure 1. Monthly rain fall distribution of the research sites during the study (2013).
Source: National Meteorological Agency.

best produce the highest MRR at all locations.

Conclusion

This study was aimed to develop improved agronomic package for malt barley production for producers in

Tigray region in response to the newly constructed brewery factory (Raya Beer). Therefore, based on the performance in grain yield, malt quality traits and net profit and assuming similar environmental conditions, the interaction of variety Behatty at 100 kg ha⁻¹ seed rate and 46/69 kg ha⁻¹ N/P₂O₅ was the best agronomic package at




Figure 2. Effect of N-fertilizer rate on grain protein content, HLW, TKW and % of plump kernels of malt barley.

Table 8. Interaction effect of N-rate, seed rate and variety on grain protein across location.

N-rate kg ha ⁻¹	Ofla				Enda- Mekoni				Emba-Alaje			
	100 kg ha ⁻¹		150 kg ha ⁻¹		100 kg ha ⁻¹		150 kg ha ⁻¹		100 kg ha ⁻¹		150 kg ha ⁻¹	
	Bekoji-1	Bhatt y	Bekoji-1	Bhatt y	Bekoji-1	Bhatt y	Bekoji-1	Bhatt y	Bekoji-1	Bhatt y	Bekoji-1	Bhatt y
0	11.17	9.48	10.43	9.53	10.60	10.03	10.19	9.75	10.19	9.73	10.073	9.69
23	11.57	9.87	11.09	9.90	11.40	10.72	10.82	10.50	11.102	10.71 2	10.776	10.24 3
34.5	11.98	10.34	11.40	9.99	11.69	10.91	11.00	10.60	11.469	10.87 5	11.394	10.67 1
46	12.19	10.79	11.89	10.23	11.78	11.16	11.32	11.10	11.5	10.89 9	11.37	10.58 7
57.5	12.32	11.01	12.14	11.14	12.03	11.57	11.66	11.03	11.55	11.28 7	12.073	11.19 3
69	12.50	11.22	12.26	11.57	12.81	12.09	12.69	11.66	11.948	11.44 3	12.136	11.68 7
80.5	12.72	11.88	12.44	11.72	13.00	12.28	12.75	11.78	12.25	11.56 2	12.261	11.73 7
Mean	12.21	10.85	11.87	10.76	12.12	11.46	11.71	11.11	11.64	11.13	11.67	11.02

Table 9. Partial budget analysis of N- fertilizer rates on malt barley production.

N-rate (kg ha ⁻¹)	Oflla			E/Mekoni			E/Alaje		
	TVC	NB	MRR	TVC	NB	MRR	TVC	NB	MRR
0	0	19957	-	0	20801.3	-	0	35445.2	-
23	750	23437	463.95	750	23829.8	403.8	750	42646.2	960.1
34.5	1125	2546	539.5	1125	26636.7	748.5	1125	44914.7	604.9
46	1475	28924	989.95	1475	29948.4	946.2	1475	47894.5	851.3
57.5	1825	29697	x	1825	31092	x	1825	49379	x
69	2175	32557	817.07	2175	34271.2	908.4	2175	53338.9	1131.4
80.5	2525	31956D	x	2525	34470.8	x	2525	50783.75D	x

TVC = total variable cost (Birr ha⁻¹), NB = Net benefit per hectare (Birr ha⁻¹), MRR = Marginal rate of return (%).

all locations. In addition, combinations of Bekoji-1 at 100 kg ha⁻¹ seed rate and 46/69 kg ha⁻¹ N/P₂O₅ can be recommended for Oflla and Emba-Alaje districts.

ACKNOWLEDGEMENTS

This research was the outcome of many individuals and institutions. First, the authors would like to thank Tigray Agricultural Research Institute (TARI) and CASCAPE-Mekele University project for financial fund support. Their acknowledgment is also extended to Professor Fassil Kebede and Dr. Yemane Tsehayu for their valuable suggestions, comments and material supports of this research.

REFERENCES

- Ahmad N (1999). Fertilizer scenario in Pakistan: policies and development. In: Proc. Conference on agriculture fertilizer use (2010). NFDC, Publication 2/99. Islamabad.
- Alam M, Haider S, Paul N (2005). Effects of sowing time and nitrogen fertilizer on barley (*Hordeum vulgare* L.) Bangladesh J. Bot. **34**(1): 27-30.
- Agricultural transformation agency (ATA) (2012). The business case for investing in a malting plant in Ethiopia.
- Bayeh M, Birhane L (2006). Barley research and development in Ethiopia: an overview In: Mulatu B and Grando S (eds). 2011. Barley Research and Development in Ethiopia. Proceedings of the 2nd National Barley Research and Development Review Workshop. 28-30 November 2006, HARC, Holetta, Ethiopia. ICARDA, Aleppo, Syria. Pp xiv + 391.
- CIMMYT (1988). From agronomic data to farmer recommendations: An Economics Workbook. Mexico, D.F.
- European brewery convention (EBC) (1987). Analytical-European Brewery Convention. Fourth Edition. Brauerei-und Getranke-Rundschau, pp-39.
- Ethiopian standard agency (ESA) (2001). Malt barley and Barley Malt- Determination of protein and soluble nitrogen content of wort. Ethiopian Standard agency, method ES 669-2001.
- Genstat 16.1.0. 2013:Genstat statistical software version 16.
- Grant D (2000). Nitrogen fertilization of dry land malt barley for yield and quality: Fertilizer Facts Number 24: December 2000: Western Triangle Ag. Research Center, Conrad. Access online. http://landresources.montana.edu/fertilizerfacts/24_Nitrogen_Fertilization_of_Dryland_malt_Barley.htm
- Landon JR (Ed.) 1991. Booker tropical soil manual: a handbook for soil survey and agricultural land evaluation in the tropics. Longman Scientific & Technical, Harlow, England.
- McKenzie R, Jackson G (2005) Barley Production in Semiarid Regions-Making the Malting Grade. Better Crops 89 (4).
- Miao Y, Mulla D, Robert C and Hernandez J (2006). within field variation in corn yield and grain quality responses to nitrogen fertilization and hybrid selection. Agro. J., 98: 129-140.
- Shafi M, Bakht J, Jalal F, Aman MK, Khattak SG (2011). Effect Of Nitrogen Application On Yield And Yield Components Of Barley (*Hordeum vulgare* L.). Pak. J. Bot., 43(3): 1471-1475
- Moreno A, Moreno M, Ribas F, Cabello M (2003). Influence of nitrogen fertilizer on grain yield of barley (*Hordeum vulgare* L.) under irrigated condition. Spanish journal of agricultural Research (2003)1(1): 91-100.
- Podsiadlo C, Koszanski Z, Zbiec I (1999): Response of some spring barley cultivars to irrigation and mineral fertilization. Part II. Canopyarchitecture and yield structure. Folia Univ. Agril. Stetinensis, Agric., 73: 131-137.
- O'Donovan J, Turkington T, Edney E, Smith M, Harker K, Beres B, Clayton G, Brandt S, Johnson E, Lafond G, May W, Grant C, McKenzie R, Patricia P, Kadeta Y, Stonehouse K, Chapman B (2013). Farmtech 2013. Conference Proceedings. Access online www.farmtechconference.com (May 2014).
- O'Donovan J, Turkington T, Edney E, Clayton G, McKenzie R, Patricia PE, Lafond GP, Granth CA,

- Brandt S, Harkerj KN, Johnson kEN, May WE (2011). Seed rate, nitrogen rate, and cultivar effects on malting barley. *Crop Agro. J.* 103(3):709-716.
- Oikeh S, Chude V, Kling G, Horst W (2007). Comparative productivity of nitrogen-use efficient and nitrogen inefficient maize cultivars and traditional grain sorghum in the moist Savanna of West Africa. *African J. Agri. Res.*, 2: 112-118.
- Shafi M, Bakht J, Jalal F, Khan A, Khattak G (2011). Effect of nitrogen application on yield and yield components of barley (*Hordeum vulgare* L.). *Pak. J. Bot.*, 43(3): 1471-1475.
- Stewart W (2002). Fertilizer contributions to crop yield. In: news and views 2002. A regional newsletter published by the Potash and Phosphate Institute (PPI) and the Potash & Phosphate Institute of Canada (PPIC). Atlanta, GA 30329.
- Stougaard RN, Xue Q (2004). Spring wheat seed size and seed rate effects on yield loss due to wild oat (*Avena fatua*) interference. *Weed Sci.* 52:133–141.
- Wheat. In: *Cereal Grain Crops*. Reston Publishing Company, Reston, VA.
- Tefera A (2012). Ethiopia grain and feed annual report, gain report number: ET 1201